

PROPERTIES AND RECOMMENDED CONDITIONS FOR LONG-TERM STORAGE OF FRESH FRUITS AND VEGETABLES

Listed alphabetically according to common name

Compiled by Marita Cantwell (email: micantwell@ucdavis.edu)

Common name	Scientific name	Storage temperature		Relative humidity %	Highest freezing temperature		Ethylene* production	Ethylene** sensitivity	Approximate storage-life	Observations and beneficial CA conditions
		°C	°F		°C	°F				
Acerola; Barbados cherry	<i>Malpighia glabra</i>	0	32	85-90	-1.4	29.4			6-8 weeks	
African horned melon; Kiwano	<i>Cucumis metuliferus</i>	13-15	55-59	90			L	M	3-6 months	
Amaranth; Pigweed	<i>Amaranthus spp.</i>	0-2	32-36	95-100			VL	M	10-14 days	
Anise; Fennel	<i>Foeniculum vulgare</i>	0-2	32-36	90-95	-1.1	30.0			2-3 weeks	
Apple	<i>Malus pumila</i>									2-3% O ₂ + 1-2% CO ₂
not chilling sensitive		-1.1-0	30-32	90-95	-1.5	29.3	VH	H	3-6 months	
chilling sensitive	Yellow Newtown, Grimes Golden, McIntosh,	4	40	90-95	-1.5	29.3	VH	H	1-2 months	
Apricot	<i>Prunus armeniaca</i>	-0.5-0	31-32	90-95	-1.1	30.0	M	M	1-3 weeks	2-3% O ₂ + 2-3% CO ₂
Artichoke										
Globe artichoke	<i>Cynara scolymus</i>	0	32	95-100	-1.2	29.9	VL	L	2-3 weeks	2-3% O ₂ + 3-5% CO ₂
Chinese artichoke	<i>Stachys affinia</i>	0	32	90-95			VL	VL	1-2 weeks	
Jerusalem artichoke	<i>Helianthus tuberosus</i>	-0.5-0	31-32	90-95	-2.5	27.5	VL	L	4 months	
Arugula	<i>Eruca vesicaria var. sativa</i>	0	32	95-100			VL	H	7-10 days	
Asian pear, Nashi	<i>Pyrus serotina;</i> <i>P. pyrifolia</i>	1	34	90-95	-1.6	29.1	H	H	4-6 months	
Asparagus, green, white	<i>Asparagus officinalis</i>	2.5	36	95-100	-0.6	31.0	VL	M	2-3 weeks	5-12% CO ₂ in air
Atemoya	<i>Annona squamosa x A. cherimola</i>	13	55	85-90			H	H	2-4 weeks	3-5% O ₂ + 5-10% CO ₂
Avocado	<i>Persea americana</i>									
cv Fuerte, Hass		3-7	37-45	85-90	-1.6	29.1	H	H	2-4 weeks	2-5% O ₂ + 3-10% CO ₂
cv. Fuchs, Pollock		13	55	85-90	-0.9	30.4	H	H	2 weeks	
cv. Lula, Booth		4	40	90-95	-0.9	30.4	H	H	4-8 weeks	
Babaco, Mt. papaya	<i>Carica candamarcensis</i>	7	45	85-90					1-3 weeks	
Banana	<i>Musa paradisiaca var. sapientum</i>	13-15	55-59	90-95	-0.8	30.6	M	H	1-4 weeks	2-5% O ₂ + 2-5% CO ₂
Barbados cherry	see Acerola									
Beans										
Snap; Wax; Green	<i>Phaseolus vulgaris</i>	4-7	40-45	95	-0.7	30.7	L	M	7-10 days	2-3% O ₂ + 4-7% CO ₂
Fava, Broad beans	<i>Vicia faba</i>	0	32	90-95					1-2 weeks	
Lima beans	<i>Phaseolus lunatus</i>	5-6	41-43	95	-0.6	31.0	L	M	5-7 days	
Winged bean	<i>Psophocarpus tetragonolobus</i>	10	50	90					4 weeks	
Long bean; Yard-long	<i>Vigna sesquipedalis</i>	4-7	40-45	90-95			L	M	7-10 days	

Common name	Scientific name	Storage temperature		Relative humidity %	Highest freezing temperature		Ethylene* production	Ethylene** sensitivity	Approximate storage-life	Observations and beneficial CA conditions
		°C	°F		°C	°F				
Beet, bunched	<i>Beta vulgaris</i>	0	32	98-100	-0.4	31.3	VL	L	10-14 days	
beet, topped		0	32	98-100	-0.9	30.3	VL	L	4 months	
Berries										
Blackberries	<i>Rubus spp.</i>	-0.5-0	31-32	90-95	-0.8	30.6	L	L	3-6 days	5-10% O2 + 15-20% CO2
Blueberries	<i>Vaccinium corymbosum</i>	-0.5-0	31-32	90-95	-1.3	29.7	L	L	10-18 days	2-5% O2 + 12-20% CO2
Cranberry	<i>Vaccinium macrocarpon</i>	2-5	35-41	90-95	-0.9	30.4	L	L	8-16 weeks	1-2% O2 + 0-5% CO2
Dewberry	<i>Rubus spp.</i>	-0.5-0	31-32	90-95	-1.3	29.7	L	L	2-3 days	
Elderberry	<i>Rubus spp.</i>	-0.5-0	31-32	90-95	-1.1	30.0	L	L	5-14 days	
Loganberry	<i>Rubus spp.</i>	-0.5-0	31-32	90-95	-1.7	28.9	L	L	2-3 days	
Raspberries	<i>Rubus idaeus</i>	-0.5-0	31-32	90-95	-0.9	30.4	L	L	3-6 days	5-10% O2 + 15-20% CO2
Strawberry	<i>Fragaria spp.</i>	0	32	90-95	-0.8	30.6	L	L	7-10 days	5-10% O2 + 15-20% CO2
Bittermelon; Bitter gourd	<i>Momordica charantia</i>	10-12	50-54	85-90			L	M	2-3 weeks	2-3%O2 + 5% CO2
Black salsify; Scorzonera	<i>Scorzonera hispanica</i>	0-1	32-34	95-98			VL	L	6 months	
Bok choy	<i>Brassica chinensis</i>	0	32	95-100			VL	H	3 weeks	
Breadfruit	<i>Artocarpus altilis</i>	13-15	55-59	85-90					2-4 weeks	
Broccoli	<i>B. oleracea</i> var. <i>Italica</i>	0	32	95-100	-0.6	31.0	VL	H	10-14 days	1-2% O2 + 5-10% CO2
Brussels sprouts	<i>Brassica oleracea</i> var. <i>Gemnjfera</i>	0	32	95-100	-0.8	30.5	VL	H	3-5 weeks	1-2% O2 + 5-7% CO2
Cabbage										
Chinese; Napa	<i>Brassica campestris</i> var. <i>Pekinensis</i>	0	32	95-100	-0.9	30.4	VL	M-H	2-3 months	1-2% O2 + 0-5% CO2
Common, early crop	<i>B. oleracea</i> var. <i>Capitata</i>	0	32	98-100	-0.9	30.4	VL	H	3-6 weeks	
late crop	“	0	32	95-100	-0.9	30.4	VL	H	5-6 months	3-5% O2 + 3-7% CO2
Cactus pads or stems, Nopalitos	<i>Opuntia spp.</i>	5-10	41-50	90-95			VL	M	2-3 weeks	
Cactus fruit; Prickly pear fruit	<i>Opuntia spp.</i>	5	41	85-90	-1.8	28.7	VL	M	2-6 weeks	2%O2 +2-5%CO2
Caimito	see Sapotes									
Calamondin	see Citrus									
Canistel	see Sapotes									
Carambola, Starfruit	<i>Averrhoa carambola</i>	9-10	48-50	85-90	-1.2	29.8			3-4 weeks	
Carrots, topped	<i>Daucus carota</i>	0	32	98-100	-1.4	29.5	VL	H	3-6 months	no CA benefit; ethylene causes bitterness
bunched; immature	“	0	32	98-100	-1.4	29.5	VL	H	10-14 days	ethylene causes bitterness
Cashew apple	<i>Anacardium occidentale</i>	0-2	32-36	85-90					5 weeks	
Cassava, Yucca, Manioc	<i>Manihot esculenta</i>	0-5	32-41	85-90			VL	L	1-2 months	no CA benefit
Cauliflower	<i>Brassica oleracea</i> var. <i>Botrytis</i>	0	32	95-98	-0.8	30.6	VL	H	3-4 weeks	2-5% O2 + 2-5% CO2

Common name	Scientific name	Storage temperature		Relative humidity %	Highest freezing temperature		Ethylene* production	Ethylene** sensitivity	Approximate storage-life	Observations and beneficial CA conditions
		°C	°F		°C	°F				
Celeriac	<i>Apium graveolens</i> var. <i>Rapaceum</i>	0	32	98-100	-0.9	30.4	VL	L	6-8 months	2-4%O2 + 2-3%CO2
Celery	<i>Apium graveolens</i> var. <i>Dulce</i>	0	32	98-100	-0.5	31.1	VL	M	1-2 months	1-4%O2 + 3-5%CO2
Chard	<i>Beta vulgaris</i> var. <i>Cicla</i>	0	32	95-100			VL	H	10-14 days	
Chayote	<i>Sechium edule</i>	7	45	85-90					4-6 weeks	
Cherimoya; Custard apple	<i>Annona cherimola</i>	13	55	90-95	-2.2	28.0	H	H	2-4 weeks	3-5% O2 + 5-10% CO2
Cherries, sour	<i>Prunus cerasus</i>	0	32	90-95	-1.7	29.0	VL	L	3-7 days	3-10% O2 + 10-12% CO2
Cherries, sweet	<i>Prunus avium</i>	-1 to 0	30-32	90-95	-2.1	28.2	VL	L	2-3 weeks	10-20% O2 + 20-25% CO2
Chicory	see Endive									
Chiles	see Pepper									
Chinese broccoli; Gailan	<i>Brassica alboglabra</i>	0	32	95-100			VL	H	10-14 days	
Chinese date	See Jujube									
Chives	<i>Allium schoenoprasum</i>	0	32	95-100			VL	H	2-3 weeks	5-10%O2 + 5-10% CO2
Cilantro, Chinese parsley	See Herbs									
Citrus										
Calamondin orange	<i>Citrus reticulata</i> x <i>Fortunella</i> spp.	9-10	48-50	90	-2.0	28.3			2 weeks	
Grapefruit	<i>Citrus paradisi</i>									3-10% O2 + 5-10% CO2
CA, AZ, dry areas		14-15	58-59	85-90	-1.1	30.0	VL	M	6-8 weeks	
FL, humid areas		10-15	50-59	85-90	-1.1	30.0	VL	M	6-8 weeks	
Kumquat	<i>Fortunella japonica</i>	4	40	90-95			VL	M	2-4 weeks	
Lemon	<i>Citrus limon</i>	10-13	50-55	85-90	-1.4	29.4			1-6 months	5-10%O2 + 0-10%CO2
Lime, Mexican, Tahiti or Persian	<i>Citrus aurantifolia</i> ; <i>C. latifolia</i>	9-10	48-50	85-90	-1.6	29.1			6-8 weeks	5-10%O2 + 0-10%CO2
Orange	<i>Citrus sinensis</i>									5-10% O2 + 0-5% CO2
CA, AZ, dry areas		3-9	38-48	85-90	-0.8	30.6	VL	M	3-8 weeks	
FL; humid regions		0-2	32-36	85-90	-0.8	30.6	VL	M	8-12 weeks	
Blood orange		4-7	40-44	90-95	-0.8	30.6			3-8 weeks	
Seville; Sour	<i>Citrus aurantium</i>	10	50	85-90	-0.8	30.6	L	M	12 weeks	
Pummelo	<i>Citrus grandis</i>	7-9	45-48	85-90	-1.6	29.1			12 weeks	
Tangelo, Minneola	<i>C. reticulata</i> x <i>paradisi</i>	7-10	45-50	85-95	-0.9	30.3			2-4 weeks	
Tangerine, Mandarin	<i>Citrus reticulata</i>	4-7	40-45	90-95	-1.1	30.1	VL	M	2-4 weeks	
Coconut	<i>Cocos nucifera</i>	0-2	32-36	80-85	-0.9	30.4			1-2 months	
Collards	<i>B. oleracea</i> var. <i>Acephala</i>	0	32	95-100	-0.5	31.1	VL	H	10-14 days	
Corn, sweet and baby	<i>Zea mays</i>	0	32	95-98	-0.6	30.9	VL	L	5-8 days	2-4%O2 + 5-10%CO2; to 4 wks, 5-10%O2+15%CO2
Cowpeas	See Peas									
Cucumber, slicing	<i>Cucumis sativus</i>	10-12	50-54	85-90	-0.5	31.1	L	H	10-14 days	3-5% O2 + 0-5% CO2
pickling		4	40	95-100			L	H	7 days	3-5% O2 + 3-5% CO2

Common name	Scientific name	Storage temperature		Relative humidity %	Highest freezing temperature		Ethylene* production	Ethylene** sensitivity	Approximate storage-life	Observations and beneficial CA conditions
		°C	°F		°C	°F				
Currants	<i>Ribes sativum</i> ; <i>R. nigrum</i> ; <i>R. rubrum</i>	-0.5-0	31-32	90-95	-1.0	30.2	L	L	1-4 weeks	CA can extend storage life to 3-6 months
Custard apple	see Cherimoya									
Daikon; Oriental radish	<i>Raphanus sativus</i>	0-1	32-34	95-100			VL	L	4 months	
Dasheen	see Taro									
Date	<i>Phoenix dactylifera</i>	-18-0	0-32	75	-15.7	3.7	VL	L	6-12 months	
Dill	see Herbs									
Durian	<i>Durio zibethinus</i>	4-6	39-42	85-90					6-8 weeks	3-5% O2 + 5-15% CO2
Eggplant	<i>Solanum melongena</i>	10-12	50-54	90-95	-0.8	30.6	L	M	1-2 weeks	3-5% O2 + 0% CO2
Endive, Escarole	<i>Cichorium endivia</i>	0	32	95-100	-0.1	31.7	VL	M	2-4 weeks	
Belgian endive; Witloof chicory	<i>Cichorium intybus</i>	2-3	36-38	95-98			VL	M	2-4 weeks	light causes greening; 3-4%O2 + 4-5%CO2
Epazote	See Herbs									
Fava bean	See Beans									
Feijoa, Pineapple guava	<i>Feijoa sellowiana</i>	5-10	41-50	90			M	L	2-3 weeks	
Fennel, see anise										
Fig	<i>Ficus carica</i>	-0.5-0	31-32	85-90	-2.4	27.6	M	L	7-10 days	5-10%O2 + 15-20%CO2
Garlic bulb	<i>Allium sativum</i>	-1-0	30-32	65-70	-2.0	28.4	VL	L	6-7 months	0.5%O2 + 5-10%CO2
Ginger	<i>Zingiber officinale</i>	13	55	65			VL	L	6 months	no CA benefit
Gooseberry	<i>Ribes grossularia</i>	-0.5-0	31-32	90-95	-1.1	30.0	L	L	3-4 weeks	
Granadilla	see Passionfruit									
Grape	<i>Vitis vinifera</i> <i>a=fruit; b=stem</i>	-0.5 - 0	31-32	90-95	-2.7 a -2.0 b	27.1 a 28.4 b	VL	L	1-6 months	2-5%O2 + 1-3%CO2; to 4 wks, 5-10%O2+10-15%CO2
American grape	<i>Vitis labrusca</i>	-1 to -0.5	30-31	90-95	-1.4	29.4	VL	L	2-8 weeks	
Grapefruit	see Citrus									
Guava	<i>Psidium guajava</i>	5-10	41-50	90			L	M	2-3 weeks	
Herbs, fresh culinary	See specific herb									5-10%O2 + 5-10%CO2
Basil	<i>Ocimum basilicum</i>	10	50	90			VL	H	7 days	2%O2 + 0 to <10%CO2
Chives	<i>Allium schoenoprasum</i>	0	32	95-100	-0.9	30.4	L	M		
Cilantro, Chinese parsley	<i>Coriandrum sativum</i>	0-1	32-34	95-100			VL	H	2 weeks	3%O2 + 7-10%CO2; air + 7-10%CO2
Dill	<i>Anethum graveolens</i>	0	32	95-100	-0.7	30.7	VL	H	1-2 weeks	5-10%O2 + 5-10% CO2
Epazote	<i>Chenopodium ambrosioides</i>	0-5	32-41	90-95			VL	M	1-2 weeks	
Mint	<i>Mentha spp.</i>	0	32	95-100			VL	H	2-3 weeks	5-10%O2 + 5-10% CO2
Oregano	<i>Origanum vulgare</i>	0-5	32-41	90-95			VL	M	1-2 weeks	
Parsley	<i>Petroselinum crispum</i>	0	32	95-100	-1.1	30.0	VL	H	1-2 months	5-10%O2 + 5-10% CO2
Perilla, Shiso	<i>Perilla frutescens</i>	10	50	95			VL	M	7 days	
Sage	<i>Salvia officinalis</i>	0	32	90-95					2-3 weeks	
Thyme	<i>Thymus vulgaris</i>	0	32	90-95					2-3 weeks	
Horseradish	<i>Armoracia rusticana</i>	-1 to 0	30-32	98-100	-1.8	28.7	VL	L	10-12 mo.	

Common name	Scientific name	Storage temperature		Relative humidity %	Highest freezing temperature		Ethylene* production	Ethylene** sensitivity	Approximate storage-life	Observations and beneficial CA conditions
		°C	°F		°C	°F				
Husk tomato	see Tomatillo									
Jaboticaba	<i>Myrciaria cauliflora</i> = <i>Eugenia cauliflora</i>	13-15	55-59	90-95					2-3 days	
Jackfruit	<i>Artocarpus heterophyllus</i>	13	55	85-90			M	M	2-4 weeks	
Jerusalem artichoke	see Artichoke									
Jicama, Yambean	<i>Pachyrrhizus erosus</i>	13-18	55-65	85-90			VL	L	1-2 months	
Jujube; Chinese date	<i>Ziziphus jujuba</i>	2.5-10	36-50	85-90	-1.6	29.2	L	M	1 month	
Kaki	see Persimmon									
Kale	<i>Brassica oleracea</i> var. <i>acephala</i>	0	32	95-100	-0.5	31.1	VL	H	10-14 days	
Kiwano	see African horned melon									
Kiwifruit; Chinese gooseberry	<i>Actinidia chinensis</i>	0	32	90-95	-0.9	30.4	L	H	3-5 months	1-2% O2 + 3-5% CO2
Kohlrabi	<i>Brassica oleracea</i> var. <i>Gongylodes</i>	0	32	98-100	-1.0	30.2	VL	L	2-3 months	no CA benefit
Kumquat	See Citrus									
Langsat; Lanzone	<i>Aglaia</i> sp.; <i>Lansium</i> sp.	11-14	52-58	85-90					2 weeks	
Leafy Greens										
Cool season	<i>various genera</i>	0	32	95-100	-0.6	31.0	VL	H	10-14 days	
Warm season	<i>various genera</i>	7-10	45-50	95-100	-0.6	31.0	VL	H	5-7 days	
Leek	<i>Allium porrum</i>	0	32	95-100	-0.7	30.7	VL	M	2 months	1-2% O2 +2-5% CO2
Lemon	see Citrus									
Lettuce	<i>Lactuca sativa</i>	0	32	98-100	-0.2	31.7	VL	H	2-3 weeks	2-5% O2 + 0% CO2
Lima bean	see Beans									
Lime	see Citrus									
LoBok	see Daikon									
Longan	<i>Dimocarpus longan</i> = <i>Euphoria longan</i>	4-7	39-45	90-95	-2.4	27.7			2-4 weeks	
Long bean	See Beans									
Loquat	<i>Eriobotrya japonica</i>	0	32	90-95	-1.9	28.6			3 weeks	
Luffa; Chinese okra	<i>Luffa</i> spp.	10-12	50-54	90-95			L	M	1-2 weeks	
Lychee, Litchi	<i>Litchi chinensis</i>	1-2	34-36	90-95			M	M	3-5 weeks	3-5% O2 + 3-5% CO2
Malanga, Tania, New cocoyam	<i>Xanthosoma</i> <i>sagittifolium</i>	7	45	70-80			VL	L	3 months	
Mamey	see Sapote									
Mandarin	see Citrus									
Mango	<i>Mangifera indica</i>	13	55	85-90	-1.4	29.5	M	M	2-3 weeks	3-5%O2 + 5-10%CO2
Mangosteen	<i>Garcinia mangostana</i>	13	55	85-90			M	H	2-4 weeks	
Melons										
Cantaloupes and other netted melons	<i>Cucurbita melo</i> var. <i>reticulatus</i>	2-5	36-41	95	-1.2	29.9	H	M	2-3 weeks	3-5% O2 + 10-15% CO2

Common name	Scientific name	Storage temperature		Relative humidity %	Highest freezing temperature		Ethylene* production	Ethylene** sensitivity	Approximate storage-life	Observations and beneficial CA conditions
		°C	°F		°C	°F				
Casaba	<i>Cucurbita melo</i>	7-10	45-50	85-90	-1.0	30.3	L	L	3-4 weeks	3-5% O2 + 5-10% CO2
Crenshaw	<i>Cucurbita melo</i>	7-10	45-50	85-90	-1.1	30.1	M	H	2-3 weeks	3-5% O2 + 5-10% CO2
Honeydew, and Orange-flesh	<i>Cucurbita melo</i>	5-10	41-50	85-90	-1.1	30.1	M	H	3-4 weeks	3-5% O2 + 5-10% CO2
Persian	<i>Cucurbita melo</i>	7-10	45-50	85-90	-0.8	30.6	M	H	2-3 weeks	3-5% O2 + 5-10% CO2
Mint	see Herbs									
Mombin	see Spondias									
Mushrooms	<i>Agaricus, other genera</i>	0	32	90	-0.9	30.4	VL	M	7-14 days	3-21%O2 + 5-15%CO2
Mustard greens	<i>Brassica juncea</i>	0	32	90-95			VL	H	7-14 days	
Nashi	see Asian pear									
Nectarine	<i>Prunus persica</i>	-0.5-0	31-32	90-95	-0.9	30.3	M	M	2-4 weeks	1-2% O2 + 3-5% CO2; internal breakdown 3-10°C
Okra	<i>Abelmoschus esculentus</i>	7-10	45-50	90-95	-1.8	28.7	L	M	7-10 days	air + 4-10%CO2
Olives, fresh	<i>Olea europea</i>	5-10	41-50	85-90	-1.4	29.4	L	M	4-6 weeks	2-3% O2 + 0-1% CO2
Onions	<i>Allium cepa</i>									
Mature bulbs, dry		0	32	65-70	-0.8	30.6	VL	L	1-8 months	1-3% O2 + 5-10% CO2
Green onions		0	32	95-100	-0.9	30.4	L	H	3 weeks	2-4% O2 + 10-20% CO2
Orange	see Citrus									
Papaya	<i>Carica papaya</i>	7-13	45-55	85-90	-0.9	30.4	M	M	1-3 weeks	2-5% O2 + 5-8% CO2
Parsley	see Herbs									
Parsnips	<i>Pastinaca sativa</i>	0	32	95-100	-0.9	30.4	VL	H	4-6 months	ethylene causes bitterness
Passionfruit	<i>Passiflora spp.</i>	10	50	85-90			VH	M	3-4 weeks	
Peach	<i>Prunus persica</i>	-0.5-0	31-32	90-95	-0.9	30.3	M	M	2-4 weeks	1-2%O2 + 3-5%CO2; internal breakdown 3-10°C
Pear, European	<i>Pyrus communis</i>	-1.5 to -0.5	29-31	90-95	-1.7	29.0	H	H	2-7 months	Cultivar variations; 1-3%O2 + 0-5% CO2
Pear, Asian	See Asian Pear									
Peas in pods; Snow, Snap & Sugar peas	<i>Pisum sativum</i>	0	32	90-98	-0.6	30.9	VL	M	1-2 weeks	2-3% O2 + 2-3% CO2
Southern peas; Cowpeas	<i>Vigna sinensis</i> = <i>V. unguiculata</i>	4-5	40-41	95					6-8 days	
Pepino; Melon pear	<i>Solanum muricatum</i>	5-10	41-50	95			L	M	4 weeks	
Peppers										
Bell Pepper, Paprika	<i>Capsicum annuum</i>	7-10	45-50	95-98	-0.7	30.7	L	L	2-3 weeks	2-5% O2 + 2-5% CO2
Hot peppers, Chiles	<i>Capsicum annuum and C. frutescens</i>	5-10	41-50	85-95	-0.7	30.7	L	M	2-3 weeks	3-5% O2 + 5-10% CO2
Persimmon; Kaki	<i>Diospyros kaki</i>									3-5% O2 + 5-8% CO2
Fuyu		0	32	90-95	-2.2	28.0	L	H	1-3 months	
Hachiya		0	32	90-95	-2.2	28.0	L	H	2-3 months	
Pineapple	<i>Ananas comosus</i>	7-13	45-55	85-90	-1.1	30.0	L	L	2-4 weeks	2-5% O2 + 5-10% CO2
Plantain	<i>Musa paradisiaca var. paradisiaca</i>	13-15	55-59	90-95	-0.8	30.6	L	H	1-5 weeks	

Common name	Scientific name	Storage temperature		Relative humidity %	Highest freezing temperature		Ethylene* production	Ethylene** sensitivity	Approximate storage-life	Observations and beneficial CA conditions
		°C	°F		°C	°F				
Plums and Prunes	<i>Prunus domestica</i>	-0.5 - 0	31-32	90-95	-0.8	30.5	M	M	2-5 weeks	1-2%O ₂ + 0-5%CO ₂
Pomegranate	<i>Punica granatum</i>	5-7.2	41-45	90-95	-3.0	26.6	VL	L	2-3 months	3-5% O ₂ + 5-10% CO ₂
Potato, early crop	<i>Solanum tuberosum</i>	10-15	50-59	90-95	-0.8	30.5	VL	M	10-14 days	no CA benefit
late crop		4-8	40-46	95-98	-0.8	30.5	VL	M	5-10 months	no CA benefit
Pumpkin	<i>Cucurbita maxima</i>	12-15	54-59	50-70	-0.8	30.5	L	M	2-3 months	
Quince	<i>Cydonia oblonga</i>	-0.5-0	31-32	90	-2.0	28.4	L	H	2-3 months	
Radicchio	<i>Cichorium intybus</i>	0-1	32-34	95-100					4-8 weeks	
Radish	<i>Raphanus sativus</i>	0	32	95-100	-0.7	30.7	VL	L	1-2 months	1-2%O ₂ + 2-3%CO ₂
Rambutan	<i>Nephelium lappaceum</i>	12	54	90-95			H	H	1-3 weeks	3-5% O ₂ + 7-12% CO ₂
Rhubarb	<i>Rheum rhaponticum</i>	0	32	95-100	-0.9	30.3	VL	L	2-4 weeks	
Rutabaga	<i>Brassica napus</i> var. <i>Napobrassica</i>	0	32	98-100	-1.1	30.1	VL	L	4-6 months	
Sage	see Herbs									
Salsify; Vegetable oyster	<i>Trapopogon porrifolius</i>	0	32	95-98	-1.1	30.1	VL	L	2-4 months	
Sapotés										
Caimito, Star apple	<i>Chrysophyllum cainito</i>	3	38	90	-1.2	29.9			3 weeks	
Canistel, Eggfruit	<i>Pouteria campechiana</i>	13-15	55-59	85-90	-1.8	28.7			3 weeks	
Black sapote	<i>Diospyros ebenaster</i>	13-15	55-59	85-90	-2.3	27.8			2-3 weeks	
White sapote	<i>Casimiroa edulis</i>	20	68	85-90	-2.0	28.4			2-3 weeks	
Mamey sapote	<i>Calocarpum mammosum</i>	13-15	55-59	90-95			H	H	2-3 weeks	
Sapodilla, Chicosapote	<i>Achras sapota</i>	15-20	59-68	85-90			H	H	2 weeks	
Scorzonera	see Black Salsify									
Shallots	<i>Allium cepa</i> var <i>ascalonicum</i>	0-2.5	32-36	65-70	-0.7	30.7	L	L		
Snap bean	see Beans									
Soursop	<i>Annona muricata</i>	13	55	85-90					1-2 weeks	3-5%O ₂ + 5-10% CO ₂
Spinach	<i>Spinacia oleracea</i>	0	32	95-100	-0.3	31.5	VL	H	10-14 days	5-10%O ₂ + 5-10%CO ₂
Spondias, Mombin, Wi apple, Jobo, Hogplum	<i>Spondias spp.</i>	13	55	85-90					1-2 weeks	
Sprouts from seeds	<i>various genera</i>	0	32	95-100					5-9 days	
Alfalfa sprouts	<i>Medicago sativa</i>	0	32	95-100					7 days	
Bean sprouts	<i>Phaseolus sp.</i>	0	32	95-100					7-9 days	
Radish sprouts	<i>Raphanus sp.</i>	0	32	95-100					5-7 days	
Squash										
Summer (soft rind); Courgette	<i>Cucurbita pepo</i>	7-10	45-50	95	-0.5	31.1	L	M	1-2 weeks	3-5% O ₂ + 5-10% CO ₂
Winter (hard rind); Calabash	<i>Cucurbita moschata</i> ; <i>C. maxima</i>	12-15	54-59	50-70	-0.8	30.5	L	M	2-3 months	large differences among varieties
Star-apple	see Sapotes									
Starfruit	see Carambola									
Strawberry	see Berries									

Common name	Scientific name	Storage temperature		Relative humidity %	Highest freezing temperature		Ethylene* production	Ethylene** sensitivity	Approximate storage-life	Observations and beneficial CA conditions
		°C	°F		°C	°F				
Sweetpotato, "Yam"	<i>Ipomea batatas</i>	13-15	55-59	85-95	-1.3	29.7	VL	L	4-7 months	
Sweetsop; Sugar apple; Custard apple	<i>Annona squamosa</i> ; <i>Annona spp.</i>	7	45	85-90			H	H	4 weeks	3-5% O2 + 5-10% CO2
Tamarillo, Tree tomato	<i>Cyphomandra betacea</i>	3-4	37-40	85-95			L	M	10 weeks	
Tamarind	<i>Tamarindus indica</i>	2-7	36-45	90-95	-3.7	25.3	VL	VL	3-4 weeks	
Taro, Cocoyam, Eddoe, Dasheen	<i>Colocasia esculenta</i>	7-10	45-50	85-90	-0.9	30.3			4 months	No CA benefit
Thyme	see Herbs									
Tomatillo; Husk tomato	<i>Physalis ixocarpa</i>	7-13	45-55	85-90			VL	M	3 weeks	
Tomato, mature-green	<i>Lycopersicon esculentum</i>	10-13	50-55	90-95	-0.5	31.0	VL	H	2-5 weeks	3-5%O2 + 2-3%CO2
Tomato, firm-ripe		8-10	46-50	85-90	-0.5	31.1	H	L	1-3 weeks	3-5%O2 + 3-5%CO2
Turnip root	<i>Brassica campestris</i> var. <i>Rapifera</i>	0	32	95	-1.0	30.1	VL	L	4-5 months	
Water chestnut	<i>Eleocharis dulcis</i>	1-2	32-36	85-90					2-4 months	
Watercress; Garden cress	<i>Lepidium sativum</i> ; <i>Nasturtium officinales</i>	0	32	95-100	-0.3	31.5	VL	H	2-3 weeks	
Watermelon	<i>Citrullus vulgaris</i>	10-15	50-59	90	-0.4	31.3	VL	H	2-3 weeks	no CA benefit
Winged bean	See Beans									
Witloof chicory	See Endive									
Yam	<i>Dioscorea spp.</i>	15	59	70-80	-1.1	30.0	VL	L	2-7 months	
Yard-long bean	See Beans									
Yucca	see Cassava									

*Ethylene production rate:

VL = very low (<0.1 µL/kg-hr at 20°C)

L = low (0.1-1.0 µL/kg-hr)

M = moderate (1.0-10.0 µL/kg-hr)

H = high (10-100 µL/kg-hr)

VH = very high (>100 µL/kg-hr)

**Ethylene sensitivity (detrimental effects include yellowing, softening, increased decay, abscission or loss of leaves, browning)

L = low sensitivity

M = moderately sensitive

H = highly sensitive

Sources of information:

Facciola, S. 1990. Cornucopia. A Source Book of Edible Plants, Kampong Publ., Vista CA. 678 pp.

Hardenburg, R., A.E. Watada, C.Y. Wang. 1986. The Commercial Storage of Fruits, Vegetables, and Florist and Nursery Stocks. USDA Agric. Hdbk No. 66. 130 pp.

Kader, A.A. et al. 1992. Postharvest Technology of Horticultural Crops. Univ. Calif. Publication 3311. 296 pp.

Kays, S.J. and J.C. Silva Dias. 1996. Cultivated Vegetables of the World. Exon Press, Athens, GA. 170 pp.

McGregor, B.M. 1987. Manual de Transporte de Productos Tropicales, USDA Agric. Handbook No. 668. 148 pp.

Maersk, Sealand and APL shipping guides

Proceedings 6th and 7th International Controlled Atmosphere Research Conferences.

Rubatzky, V.E. and M. Yamaguchi. 1997. World Vegetables. Principles, Production and Nutritive Values. 2nd ed. Chapman & Hall, N.Y. 843 pp.

Whiteman, T.M. 1957. Freezing points of Fruits, Vegetables, and Florist Stocks. USDA Mkt. Res. Rpt. No. 196. 32 pp.

Unpublished data on specialty vegetables and fruits from Marita Cantwell and Adel Kader.